

Department of Industrial Engineering and Management

Technical Report

No. 2011-9

A proof by the simplex method for the diameter of a $(0, 1)$ -polytope

Tomonari Kitahara and Shinji Mizuno



August, 2011

Tokyo Institute of Technology

2-12-1 Ookayama, Meguro-ku, Tokyo 152-8552, JAPAN
<http://www.me.titech.ac.jp/index-e.html>

A Proof by the Simplex Method for the Diameter of a $(0,1)$ -Polytope

Tomonari Kitahara* and Shinji Mizuno†

August 2011

Abstract

Naddef [3] shows that the Hirsch conjecture is true for $(0,1)$ -polytopes by proving that the diameter of any $(0,1)$ -polytope in d -dimensional Euclidean space is at most d . In this short paper, we give a simple proof for the diameter. The proof is based on the number of solutions generated by the simplex method for a linear programming problem. Our work is motivated by Kitahara and Mizuno [2], in which they get upper bounds for the number of different solutions generated by the simplex method.

1 Introduction

Let d be a positive integer and P be a polytope in the d -dimensional Euclidean space \mathbb{R}^d with f facets. Let $V(P)$ be a set of its vertices. The diameter $\delta(P)$ of the polytope P is the minimum value δ with the following property: for any two vertices $\hat{\mathbf{y}}, \tilde{\mathbf{y}}$ in $V(P)$, there exists a path from $\hat{\mathbf{y}}$ to $\tilde{\mathbf{y}}$ with the length at most δ . Hirsch conjectures that $\delta(P) \leq (f - d)$ in the context of the simplex method [1]. In a celebrated paper [4], Santos presents a counterexample to the conjecture. However, the conjecture is true for some special cases. We call P a $(0, 1)$ -polytope when any element of $\mathbf{y} \in V(P)$ is 0

*Graduate School of Decision Science and Technology, Tokyo Institute of Technology, 2-12-1-W9-62, Oo-Okayama, Meguro-ku, Tokyo, 152-8552, Japan. Tel.: +81-3-5734-2896, Fax: +81-3-5734-2947, E-mail: kitahara.t.ab@m.titech.ac.jp.

†Graduate School of Decision Science and Technology, Tokyo Institute of Technology, 2-12-1-W9-58, Oo-Okayama, Meguro-ku, Tokyo, 152-8552, Japan. Tel.: +81-3-5734-2816, Fax: +81-3-5734-2947, E-mail: mizuno.s.ab@m.titech.ac.jp.

or 1. Naddef [3] shows that the Hirsch conjecture is true for $(0, 1)$ -polytopes by proving that the inequality

$$\delta(P) \leq d \tag{1}$$

holds for any $(0, 1)$ -polytope $P \subset \mathfrak{R}^d$.

In this short paper, we give a simple proof of the inequality (1) by using the number of different solutions generated by the simplex method. Our work is motivated by Kitahara and Mizuno [2], in which they get upper bounds for the number of different solutions generated by the simplex method.

2 Proof by the simplex method

Let $P \subset \mathfrak{R}^d$ be a $(0, 1)$ -polytope with f facets and $V(P)$ be a set of its vertices. It is well known that P can be expressed by using f linear inequalities like

$$P = \{\mathbf{y} \in \mathfrak{R}^d \mid \mathbf{Q}^T \mathbf{y} \leq \mathbf{r}\}$$

for some $\mathbf{Q} \in \mathfrak{R}^{d \times f}$ and $\mathbf{r} \in \mathfrak{R}^f$. For any fixed vertex $\hat{\mathbf{y}} = (\hat{y}_1, \hat{y}_2, \dots, \hat{y}_d)^T \in V(P)$, we define a linear function

$$l(\mathbf{y}; \hat{\mathbf{y}}) = \sum_{i: \hat{y}_i=0} (1 - y_i) + \sum_{i: \hat{y}_i=1} y_i, \tag{2}$$

where $\mathbf{y} = (y_1, y_2, \dots, y_d)^T$ is a vector of variables. A linear programming problem, which maximize the linear function (2) on P , is written as

$$\begin{aligned} \max \quad & l(\mathbf{y}; \hat{\mathbf{y}}), \\ \text{subject to} \quad & \mathbf{Q}^T \mathbf{y} \leq \mathbf{r}. \end{aligned} \tag{3}$$

Obviously, this problem has the unique optimal solution $\hat{\mathbf{y}}$ whose objective function value is d .

Theorem 1 *When we generate a sequence of vertices from any initial vertex $\tilde{\mathbf{y}} \in V(P)$ by the simplex method for solving (3) so that the objective function value increases whenever an iterate is updated, the number of different vertices generated is at most d .*

Proof: From the definition (2), it is easy to see that

$$0 \leq l(\mathbf{y}; \hat{\mathbf{y}}) \leq d$$

for any $\mathbf{y} \in V(P)$. Since the objective function value is integral and increases at least one whenever an iterate is updated, the number of different vertices generated is at most d . ■

From Theorem 1, we can easily get the next result.

Theorem 2 *The diameter of any $(0, 1)$ -polytope $P \subset \mathbb{R}^d$ is at most d .*

Proof: Let $\hat{\mathbf{y}}$ and $\tilde{\mathbf{y}}$ be any two vertices of P . We define the linear function (2) and the linear programming problem (3). If we generate a sequence of vertices from the initial vertex $\tilde{\mathbf{y}}$ by the simplex method with Bland's pivoting rule, we can find the optimal solution $\hat{\mathbf{y}}$ in a finite number of iterations. From Theorem 1, the number of vertices generated is at most d . Hence there exists a path from $\hat{\mathbf{y}}$ to $\tilde{\mathbf{y}}$ with the length at most d . ■

Acknowledgment

This research is supported in part by Grant-in-Aid for Young Scientists (B) 23710164 and Grant-in-Aid for Science Research (A) 20241038 of Japan Society for the Promotion of Science.

References

- [1] G. B. Dantzig, *Linear Programming and Extensions*, Princeton University Press, Princeton, New Jersey, 1963.
- [2] T. Kitahara and S. Mizuno, A bound for the number of different basic solutions generated by the simplex method. *To appear in Mathematical Programming*.
- [3] D. Naddef, The Hirsch conjecture is true for $(0,1)$ -polytopes. *Mathematical Programming* 45 (1989) 109–110.
- [4] F. Santos, A counterexample to the Hirsch conjecture. Technical paper, available at <http://arxiv.org/abs/1006.2814>, 2010.